

MA141-004 Test #3 Form A Thursday, March 21, 2019 Dr. J. Griggs

Please put all work and answers in the blue book provided. Nothing written on the test itself will be graded. Scientific calculators OK; do not use a graphing calculator. Validate all graphs with appropriate calculus techniques. Put your name, row number and seat number on the front of the blue book.

1.) Evaluate: a.) $\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2}$

b.) $\lim_{x \rightarrow 0^+} x^{x^2}$

2.) For the function $f(x) = x^3 - x^2 - x + 4$ on $[-2, 3]$, using the first and second derivative - find the critical points and points of inflection; where the function is increasing, decreasing; where it is concave up, concave down; all relative and absolute maximum and minimum values; graph the function.

3.) Use Newton's Method to find the positive root of the equation $f(x) = 2x - e^{-x} = 0$ using $x_1 = .3$, find x_2 and x_3 .

4.) A closed wooden box (top included) is to be constructed to have a volume of 45 cubic feet. The length of the box must be 5 times the height. Once it is built, it will be coated with a very expensive acrylic - so its surface area needs to be minimized. What dimensions should the box have in order to minimize the surface area?

5.) Use differentials to approximate the increase in volume of a cube if the length of each edge of the cube is changed from 10 cm to 10.1 cm. What is the exact change in volume?

6.) Integrate: a.) $\int (x^3 - 5 \sin x + \frac{4}{\sqrt[3]{x}}) dx$

b.) $\int (5e^{3x} - \frac{2}{x} + \frac{3}{1+x^2}) dx$

7.) Using $f(x) = x^2 - 4x + 2$ on the interval $[1, 4]$, verify the hypotheses of the Mean Value Theorem and find c in $[1, 4]$ guaranteed by the conclusion. Construct a graph of this scenario.

MA141 TEST #3A SOLUTIONS:

(7 questions; 14 points each)

$$1.) \quad a.) \quad \lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2} \xrightarrow{\text{L'HOP}} \lim_{x \rightarrow 0} \frac{e^x - 1}{2x} \xrightarrow{\text{L'HOP}}$$

$\begin{matrix} \nearrow 0 \\ \downarrow 0 \end{matrix}$

7pts

$$= \lim_{x \rightarrow 0} \frac{e^x}{2} = \left(\frac{1}{2} \right)$$

$$b.) \quad \lim_{x \rightarrow 0} x^{x^2} \text{ (rewrite)} \rightarrow \lim_{x \rightarrow 0^+} e^{\ln x^{x^2}}$$

7pts

$$= \lim_{x \rightarrow 0^+} e^{(x^2)(\ln x)} = e^{\boxed{\lim_{x \rightarrow 0^+} (x^2)(\ln x)}}$$

exponent only... $\lim_{x \rightarrow 0^+} (x^2)(\ln x)$

$$= \lim_{x \rightarrow 0^+} \frac{(\ln x)}{\frac{1}{x^2}} \xrightarrow{\text{L'HOP}} \lim_{x \rightarrow 0^+} \frac{\frac{1}{x}}{\frac{-2}{x^3}}$$

$\begin{matrix} \nearrow -\infty \\ \downarrow \infty \end{matrix}$

$$= \lim_{x \rightarrow 0^+} \frac{1}{x} \cdot \frac{x^3}{-2} = \lim_{x \rightarrow 0^+} \frac{x^2}{-2} = 0$$

$$\therefore e^0 = \boxed{1}$$

$$2.) \quad f(x) = x^3 - x^2 - x + 4 \quad \text{on } [-2, 3]$$

14pts

endpoints:

$$f(-2) = (-2)^3 - (-2)^2 - (-2) + 4 = -6 \rightarrow (-2, -6)$$

$$f(3) = (3)^3 - (3)^2 - (3) + 4 = 19 \rightarrow (3, 19)$$

(page 2)

$$f'(x) = 3x^2 - 2x - 1 = 0 \quad (f'(x) \text{ never undef})$$

$$(3x+1)(x-1) = 0$$


$$x = -\frac{1}{3} \quad x = 1$$

$$f(-\frac{1}{3}) = (-\frac{1}{3})^3 - (-\frac{1}{3})^2 - (-\frac{1}{3}) + 4 = \frac{113}{27} \approx 4.185$$

$$f(1) = 1^3 - 1^2 - 1 + 4 = 3$$

critical points: $(-\frac{1}{3}, 4.185)$ & $(1, 3)$ "FLAT"

$f'(x)$:



$$f''(x) = 6x - 2 = 0$$


$$6x = 2 \quad x = \frac{1}{3}$$

($f''(x)$ never undef)


$$f(\frac{1}{3}) = (\frac{1}{3})^3 - (\frac{1}{3})^2 - (\frac{1}{3}) + 4 = \frac{97}{27} \approx 3.59$$

$(\frac{1}{3}, 3.59)$ pt. of inflection

$f''(x)$:



(page 3)



rel max:
 $(3, 19) \hat{=} (-\frac{1}{3}, 4.185)$

rel min:
 $(1, 3) \hat{=} (-2, -6)$

ABS MAX: 19 $(3, 19)$
ABS MIN: -6 $(-2, -6)$

3.) $f(x) = 2x - e^{-x}$ $x_1 = .3$

14 pts

$$f'(x) = 2 + e^{-x}$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$\begin{aligned} x_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} = .3 - \frac{f(.3)}{f'(.3)} \\ &= .3 - \frac{[.6 - .7408]}{[2 + .7408]} = .3 + .0514 \\ &= \boxed{.3514} \end{aligned}$$


(page 4)

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)} = .3514 - \frac{[.7028 - .7037]}{[2 + .7037]}$$

$$x_3 = .3514 + .00033$$

$$x_3 = .35173$$

4.)
14 pts



$$V = 45$$

$$(H)(w)(5H) = 45$$

$$5H^2w = 45$$

$$w = \frac{45}{5H^2} = \frac{9}{H^2}$$

surface area
↑

$$S = 2[(H)(5H)] + 2[(w)(H)] + 2[(5H)(w)]$$

$$S = 10H^2 + 2wH + 10wH = 10H^2 + 12wH$$

$$S = 10H^2 + 12\left(\frac{9}{H^2}\right)(H) = 10H^2 + \frac{108}{H} = S$$

$$S' = 20H - \frac{108}{H^2} = 0$$

$$20H = \frac{108}{H^2}$$

$$\frac{20H}{1} = \frac{108}{H^2}$$

$$20H^3 = 108 \quad H^3 = \frac{108}{20} = \frac{54}{10} = 5.4$$

$$H = \sqrt[3]{5.4} \approx 1.7544 \approx 1.75 \text{ ft}$$

(page 5)

$$SH \approx 5(1.7544) \approx 8.77 \text{ ft}$$

$$W \approx \frac{9}{H^2} \approx 2.94 \text{ ft}$$

max or min?

$$S'' = 20 + \frac{216}{H^3} = +$$

$\therefore$ CONCL. UP $\therefore$ MIN

5.) CUBE $V = x^3$ $x = 10$ $\Delta x = .1$

14pts $dV = V'(x) \cdot \Delta x$
 $= (3x^2) \cdot \Delta x = (3 \cdot 10^2)(.1) = 30 \text{ cm}^3$
APPROX

$$\Delta V = (10.1)^3 - (10)^3 = 1030.301 - 1000$$
$$= 30.301 \text{ cm}^3$$

EXACT

6.) a.) $\int (x^3 - 5 \sin x + 4x^{-1/3}) dx$

7pts

$$= \frac{x^4}{4} - 5(-\cos x) + 4\left(\frac{x^{2/3}}{2/3}\right) + C$$
$$= \frac{x^4}{4} + 5 \cos x + 6x^{2/3} + C$$

(page 6)

7pts b.) $\int (5 \cdot e^{3x} - \frac{2}{x} + \frac{3}{1+x^2}) dx$
 $= 5(\frac{1}{3} \cdot e^{3x}) - 2 \ln|x| + 3 \tan^{-1} x + C$
 $= \frac{5}{3} e^{3x} - 2 \ln|x| + 3 \tan^{-1} x + C$

7.) $f(x) = x^2 - 4x + 2$ on $[1, 4]$

hypotheses: ① $f(x)$ contin on $[1, 4]$??
yes, it is a polynomial

② $f(x)$ differentiable on $(1, 4)$
yes, deriv. exists for all values in $(1, 4)$

$$f'(x) = 2x - 4 = 0$$

$$2x = 4$$

$$x = 2$$

Vertex: $(2, -2)$

$$f(1) = -1 \quad f(4) = 2$$

find c such that...

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$2c - 4 = \frac{f(4) - f(1)}{4 - 1}$$

$$2c - 4 = \frac{2 - (-1)}{4 - 1} = \frac{3}{3} = 1$$

$$2c - 4 = 1$$

$$2c = 5$$

$$c = \frac{5}{2}$$

$$\frac{5}{2} \in (1, 4)$$

